

Tracking Training Load and Its Implementation in Tactical Populations: A Narrative Review

Danny Maupin, DPhty, ATC,¹ Ben Schram, PhD, DPhty,² and Robin Orr, PhD²

¹Faculty of Health Sciences and Medicine Bond University, Bond Institute of Health and Sport, Bond University, Gold Coast, QLD, Australia; and ²Tactical Research Unit, Faculty of Health Sciences and Medicine Bond University, Bond University, Gold Coast, QLD, Australia

ABSTRACT

Tactical populations often participate in demanding physical training and perform strenuous workplace tasks, increasing injury risk. Mitigating injury risk is vital for maintaining trained personnel and should be a focus for tactical populations. One such method, tracking training load, has not been studied in-depth in tactical populations, despite documented effectiveness in elite sport. Most injuries to tactical personnel are overuse in nature and therefore may be prevented by optimizing training load. Although the methods used in elite sport may not be directly transferrable to tactical environments, they may be used to inform injury mitigation strategies in tactical populations.

INTRODUCTION

Tactical populations are inclusive of, but not limited to, firefighters, police officers, and military personnel. Despite intermittent sedentary occupational tasks, especially police (e.g., sitting in a patrol car (17)), tactical populations are required to perform tasks in stressful and physically demanding situations

Address correspondence to Danny Maupin, dmaupin@bond.edu.au.

(17). This requirement often occurs while these personnel are carrying external loads (ELs), such as body armor and other personnel protection equipment (72). ELs often vary and can range from 10 kg in general police units (72), to 22 kg in firefighters (12), whereas military personnel can carry average loads of 45 kg (65). The addition of these loads not only causes an increase in metabolic demand (18) but can also lead to an increased risk of injury (49). In police in particular, these loads further increase on admission to specialist units (38,70). Specialist police units, owing to the requirement to carry additional equipment, such as riot shields and more substantial body armor, can be required to carry loads ranging from around 20 kg (11) to 40 kg (44), which can cause even greater metabolic demands and higher risk of injury (19,49).

While each tactical population undergoes high physical stress that increases the risk of injury, the stresses placed on each of these tactical populations vary. Firefighters are typically faced with significant environmental hazards (73), military personnel are more likely to carry considerably heavier loads (67), and police officers are more likely to encounter resistant and uncooperative suspects (9). Furthermore, and

apart from the nature of the occupation itself, sport and physical training can be a leading source of injury in tactical populations (42). However, sport and physical conditioning are integral to many tactical populations, with appropriate physical training required to increase occupational capability either directly (increased task performance) or indirectly (improved general health) (14,72).

Given the high injury risk of tactical populations (49), it is vital that injury mitigation strategies are researched. One such strategy that has not been explored in depth is the monitoring and optimization of training load (TL), despite documented effectiveness in elite sport (26). The purpose of this narrative review is to explore the potential benefits and limitations of tracking TL in tactical populations with the intent of reducing injury risk while maintaining or increasing fitness. This review will summarize information regarding injuries in tactical populations and the potential usefulness of optimizing TLs. Information regarding TL, how it can be measured, as well as previous research examining the

KEY WORDS:

police; military; firefighters; injury prevention; overload

Training Load in Tactical Populations

relationship between TL and injuries, fitness, and performance will also be presented.

INJURIES IN TACTICAL POPULATIONS

With differing occupational and physical training demands, the location and incidence of injuries experienced can vary; however, injury types are known to be similar across these populations (53). In a study by Taylor et al. (82), it was found that firefighters suffered injuries at a rate of 177 per 1,000 full-time employees per annum. Firefighters were most likely to suffer joint sprains and muscle strains, accounting for 66.5% of all injuries, followed by contusion/crush injuries, mental disorders, and open wounds, with the most common sites of injury being the knee, lower back, shoulder, and ankle (82).

In law enforcement, a critical review (53) found injury rates ranging from 240 to 2,500 per 1,000 personnel per annum, with multiple articles reviewed reporting the most common region of injury being the upper extremity (32.95–43.42% of injuries), whereas another stated the most common site was other unspecified sites (63.41%), followed by torso and back (20.49%). This review found the most common injury type to be sprains and strains (42.36–94.59%), followed by other muscle pains and other natures of injury (53).

With regard to military, incidence rates have been reported to range from 23 incidents, for active duty personnel, to 34 incidents, for reserve personnel, per 100 person-years of active service, with “minor personal injury” being the most common for both groups (69). Their injuries were found to most commonly be overuse or stress syndrome, muscle strains, ankle sprains, and stress fractures, which typically occurred at or below the knee (43).

Injuries in these populations can have serious downstream effects, such as a temporary or permanent loss of experience and skills, or even worse, higher workforce strain due to the requirement of covering injured

personnel (35,62). Injuries to tactical populations can also result in high monetary costs to treat and, if needed, replace (e.g., temporarily or permanently) injured personnel, placing an increased financial burden on the organization (35,62). For example, the financial cost for medical care and salaries of soldiers in the U.S. Army is estimated to be about \$1.5 billion (USD) per year (62). By exploring causative factors for these injuries and implementing programs to reduce these injuries, some of these negative effects may be mitigated.

Numerous causative factors have been studied in tactical populations, such as poor metabolic fitness levels (51), high body mass index (71), history of previous injury (2), and smoking (10). In 2000, Kaufman et al. (43) discussed how TL could potentially be another causative factor, and how optimizing TL may decrease injury risk while promoting fitness in a military population. Previous research has shown that an excessive increase in physical activity is linked with an increased injury risk (2,83). In a study by Trank et al. (83), it was found that U.S. Navy Recruits who completed the highest running mileage (>25 miles [40.23 km]) also had a significantly higher risk of injury, with no significant change in 1.5 mile (2.41 km) run performance. Although the need to track TL is evident, a cursory search of the literature shows that little research has been performed to date with regard to quantifying TL in tactical populations. In fact, a systematic review found only 2 articles regarding TL tactical populations, but these articles focused mainly on increases in physical activity (20). Despite minimal information pertaining to the tactical environment, it has become a popular tool in elite sporting teams. Similar to the study by Trank et al. (83), a medium effect size ($r = 0.59$; confidence interval [CI] and p not reported) was found between higher running distances during a given month in endurance runners and number of injury days the following month (41). By implementing methods that track TL, sporting teams are able to

adjust training stimuli at an individual level and on a weekly basis to reduce injury risk and also increase performance (26). Given the significant costs associated with injuries and the increased risk of injury in tactical populations caused by higher physical loads, a means of monitoring TL and their associated effects on injury risk could prove beneficial for tactical populations (2,83).

TRAINING LOAD

The International Olympic Committee consensus statement defines TL as a cumulative amount of stress on an individual from single, or multiple, training sessions over time (79). TL is typically measured using either external methods (e.g., total distance run or speed) or internal methods (e.g., heart rate [HR] or rating of perceived exertion [RPE]) (26). EL has previously been defined as “any external stimulus applied to the athlete that is measured independently of their internal characteristics,” whereas internal load (IL) is “load measurable by assessing internal response factors within the biological system, which may be physiological, psychological, or other” (79). Tracking EL can communicate the total amount of work completed, whereas information yielded from IL typically conveys how the individual is adapting physiologically (79). Emphasis is placed on the importance of using both measures, as together they provide a more comprehensive view of the physical stress an individual is experiencing (26). For example, an individual may perform the same overall output (as measured by EL, such as distance) on 2 different days; however, their ability to respond to this output (as measured by IL, such as HR) may have changed. Potential differences between responses to training may provide insights as to whether the individual is in a state of “readiness” and able to tolerate high TL or “fatigue” and potentially at an increased risk of both injury and decreased performance (30).

Rapid changes in both EL and IL have previously been linked to higher risk of injury in the literature (26). One example shows 40% of injuries were the result of a rapid change in TL in the preceding week in Australian football (68). Factors further compounding the issue were proposed by Gabbett that athletes who overtrain or undertrain are at higher injury risks (26). These issues have led to the development of monitoring both IL and EL over the short-term (acute, typically 1 week) and long-term (chronic, 3–6 weeks) periods, and weekly changes to track overall and rapid changes in workload (26). There are various methods of measuring TL reported in the literature, each of which has benefits and limitations; the following sections will explore these various methods.

External measurements of training load. EL is measured in an attempt to define the total work completed (79). EL measurements can include distance run, tracking volume load during strength training sessions, and number of sprints completed, among others (26). Measuring EL has been aided with the advent of Global Positioning System (GPS) devices and the subsequent accessibility of these relatively low-cost devices, which allow for accurate measurement of the above variables (15). The use of GPS to measure EL is common practice in many professional sports including Australian Football League (AFL), rugby league, and soccer (15). Research has shown the effectiveness and validity of GPS to measure EL in these sports (13,22). However, there are concerns of the precision of GPS with regard to high rates of change in velocity, and although GPS technology has seen an increase in data accuracy, there are still some limitations in regard to very high speed running (>20 km/h) (40). Although there may be potential validity issues at high speed, it has been estimated that the speed of military marches while on level ground is 5.3 km/h (84), which may allow GPS devices to accurately record data. Practitioners and clinicians

may be able to use these devices effectively during training stations and while personnel are on duty and be able to record distance covered.

Another concern with GPS units is the high cost associated with them (7). Research has shown that using GPS on a cellular phone may provide an alternate way to measure distance, as the accuracy between the 2 was comparable with an approximately 3% variation identified when measuring distance and average speed (4). Of importance, this study compared the cellular devices to GPS devices with lower sampling rates and more limited accuracy; therefore, more variation may exist when comparing cellular devices to GPS devices with higher sampling rates (4). Using cellular phones or fitness watches may be an alternative tool for tactical practitioners to pursue if the acquisition of GPS units proves too costly. One key area of difference between tactical populations and athletics may be potential security concerns and GPS units. Recently, data collected by a wearable GPS unit showed the location of a secure U.S. military facility as discussed in an article published by Liz Sly in the Washington Post on January 29, 2018. Tactical personnel will need to ensure that the devices they are using are secure and will not lead to similar issues.

Apart from GPS devices to track EL, there are other, more cost-effective methods. One such method is using volume load per the NSCA's definition (number of repetitions \times EL [kg]), a convenient method because it does not require any additional equipment (29). Whereas repetition volume is the total number of repetitions performed during a workout, volume load also incorporates the weightlifted and is likely better at quantifying the total amount of work completed (29). For example, if a police officer completes 4 sets of 12 repetitions of bench press at 100 kg, their volume load will be 4,800 "work units," whereas their repetition volume will be 48. Volume

load can be used during physical training sessions, when tactical personnel are participating in strength training, to track and adjust based on previous and future loads. More complex methods that determine mechanical work during resistance exercises may be used during strength sessions; however, these require additional devices, such as force plates, accelerometers, and inertial sensors (7). Although these devices have been shown to have good reliability and validity, they are expensive methods to implement and may not be accessible in all populations (7).

Another method to track TL that may be of benefit when training large groups includes measuring total running distance or weight lifted using subjective measures (7). This method may be particularly useful in military populations on return from patrol, owing to their ability and operational requirement in tracking distances covered. In support of subjective measures, a systematic review on endurance athletes (41) or endurance runners found a medium effect size ($r = 0.59$; CI and p not reported) (54) ($r^2 = 0.36$, CI and $p = 0.001$) (8) between subjective reports of high total training distance and rates of injury or pain.

Finally, there are other methods that have been used to track TL, although these are unlikely to be used in tactical populations because of logistical concerns and time constraints (7). For example, despite time motion analysis being found to be effective in tracking fatigue in professional soccer players (58), it is unlikely that tactical populations would have the time or resources to record and perform film review of all training and occupational settings. Similarly, despite training diaries being used to track load, concern exists with regard to the accuracy of self-reporting (74), especially if trying to recall events that occurred at the beginning of long hours of training (36).

Training Load in Tactical Populations

Internal measurements of training load.

Although EL measurements attempt to measure the amount of work done, IL measurements are more focused on the individual's response to the TL (26). Numerous methods are available to track IL including HR, session RPE (sRPE), and questionnaires. Historically, HR has been used to track IL because of an almost linear relationship with $\dot{V}O_2\text{max}$ (80). This allows practitioners to quantify how each individual is responding to the imposed work load (39). Likewise, blood lactate is related to $\dot{V}O_2\text{max}$ and may be used to track TL and intensity (23). As this measure has also been linked to load carriage performance in soldiers (78), it may be a useful tool to track intensity and performance. Although these methods are accurate, they can be invasive and may require a high level of technical skill or the necessary technology to be performed, and therefore may not be feasible in a tactical population. As an example, although measuring HR at one set point is a simple task, requiring only a stopwatch, measuring and recording HR consistently over a period may require more sophisticated technology, such as HR chest straps (7). Using fitness watches to measure HR may be an alternative; however, research has shown that these devices are less accurate at higher HRs compared with chest straps (80). Incorporating the RPE scale may be useful, as it is both low cost and easy to administer, especially to larger groups (39).

The RPE scale (Figure 1) was designed by Borg (5), as an attempt to measure the level of physical strain during activity. It was proposed by Foster et al. (24) that this scale can be modified to a 0–10 scale (Figure 2) and a score given at the end of activity. This score can then be multiplied by the duration of the session for a measure of TL called session RPE or sRPE (24). For example, if an athlete completed a 60-minute weightlifting session and reported the session was 6 on the modified RPE,

the athlete would have a sRPE of 360 (60×6) arbitrary units (AUs). Previous research has shown that this method correlates well with blood lactate concentration and HR after exercise (25,75). In a study by Gabbett and Domrow (25), the correlations between sRPE and both HR and blood lactate concentration were found to be $r = 0.89$ and $r = 0.86$, respectively, showing a high correlation between measures. These results suggest sRPE, using the modified scale, can be used to accurately measure IL by tactical practitioners after a variety of activities, such as physical training sessions. Although tactical personnel do tend to perform physical activity throughout the day, previous research has confined the “session duration” to fixed training hours experienced each day (06:00–22:00), allowing for comparison of RPE scores across days (64). There was good agreement between RPE and recorded training impulse (TRIMP) scores ($R^2 = 0.57\text{--}0.77$), showing that this may be a valid method to compare RPE scores in tactical populations, although further research will be necessary (64).

TRIMP was developed by Bannister (3) as a method of quantifying a training session. This method is calculated by using training duration, maximal HR, resting HR, average HR, and a sex-dependent exponential coefficient (3). This equation was further modified by Manzi et al. (57), by introducing an individual-based exponential factor. The modification was done in attempts to better determine individual physiological response to exercise (57). The use of this equation, TRIMP_i , has been shown to be a valid method to track fitness and performance changes in long distance runners (57) as well as aerobic fitness in team sport (55). Another method of calculating TRIMP, Edward's TRIMP, multiplies the duration accumulated in 5 HR zones (zone 1 = 50–60% HRmax, zone 2 = 60–70%, zone 3 = 70–80%, zone 4 = 80–90%, and zone 5 = 90–100%) by a coefficient for each zone and then sums the result (21). This equation was used in military training and found agreement ($R^2 = 0.34\text{--}0.49$) between TRIMP and distance, suggesting this may be a valid

Ratings of perceived exertion (RPE)	Total quality recovery (TQR)
6	6
7 Very, very light	7 Very, very poor recovery
8	8
9 Very light	9 Very poor recovery
10	10
11 Fairly light	11 Poor recovery
12	12
13 Somewhat hard	13 Reasonable recovery
14	14
15 Hard	15 Good recovery
16	16
17 Very hard	17 Very good recovery
18	18
19 Very, very hard	19 Very, very good recovery
20	20

Figure 1. Borg RPE Scale and Total Quality Recovery Scale. Reproduced from Kentta and Hassmen (47).

Rating	Descriptor
0	Rest
1	Very, Very Easy
2	Easy
3	Moderate
4	Somewhat Hard
5	Hard
6	.
7	Very Hard
8	.
9	.
10	Maximal

Figure 2. Modified Borg RPE Scale. Reproduced from Foster et al. (24). RPE = rating of perceived exertion.

method to calculate IL, although it may be limited because of the variable training activities and prolonged time frames found in military populations (64). Using TRIMP may be an effective means to track IL over time in tactical populations, particularly Edward's TRIMP as an individual factor will not need to be calculated and it accounts for activity completed in various HR zones.

HR variability (HRV), the beat-to-beat fluctuation of resting HR, is another measure of IL (28). The autonomic nervous system (ANS) plays a crucial and complex role in maintaining the body's homeostasis (28). Although the ANS is involved in many different mechanisms, its ability to control HR and role in overuse injuries is a key factor when it comes to monitoring TL (28). Examining the physiology of this relationship is beyond the scope of this article, but to paraphrase, using the parasympathetic and sympathetic nervous systems, the ANS is able to increase and decrease HR, and release chemical mediators in response to injuries, including overuse injuries (31). HRV aims to indirectly measure the ANS by analyzing the change in time between heart beats during resting

HR (28). Significant changes from an individual's baseline may show that the ANS is under stress, potentially because of overtraining or the onset of an overuse injury (28). Given the many roles of the ANS, changes in HRV may be indicative of other types of stress than overuse injuries. However, this may be beneficial to those working with tactical populations given the amount of non-physical stress, such as psychological, that are experienced (60,63). Currently, research is inconclusive about the relationship between HRV and overtraining, with evidence both for and against (34,50,60), but this may be a valuable tool to measure the amount of stress, both physical and other, that are being experienced by tactical personnel.

Various wellness questionnaires, which examine factors such as sleep quantity and sleep quality, feelings of fatigue, and soreness, have also been used in an attempt to measure IL (7). One such survey, the Recovery-Stress Questionnaire (RSQ) for Athletes, was created to identify the extent of an athlete's physical and mental exhaustion and recoverability (16). This questionnaire consists of 19 subscales that discuss various topics such as general and emotional stress, burnout, physical complaints, and

physical recovery among others (16). The survey is periodically given out, with one study administering the survey 4 times during a 3-week period (45). It has previously been shown to be able to track the general parameters of training stress of athletes across multiple sports (16), although it is unknown how this scale will apply to those in tactical populations or if they will require a more specific questionnaire for their demands. However, given the high amounts of psychological stress experienced by tactical populations (63), the RSQ may provide useful information regarding not only the physical but emotional state of personnel.

Another questionnaire that attempts to quantify recovery is the Total Quality Recovery (TQR) process (46). This process is divided into 2 parts, the first being a perceived recovery scale (TQR perceived) (Figure 1). Structured like the Borg RPE Scale, this 6–20 scale attempts to measure an individual's perceived recovery at the end of the day (46). The second aspect of TQR (TQR action) is a more objective measure of recovery where individuals earn recovery points (20 points maximum) based on nutrition and hydration (10 points maximum), sleep and rest (4 points), and relaxation and emotional support (3 points) (Figure 3) (46,47). As this method combines both perceived and an objective score for recovery, it may be useful in tactical populations, although future research will be needed to validate this process.

Finally, although the above methods may be more complex, previous research in military populations has examined muscle soreness and physical fatigue using 0–10 scales (64). Although this study did not examine these measures in relation to injury, these simple questions may be able to track how tactical personnel are managing their current TL (64). Various questionnaires exist that track both the physical and psychological stress and recovery experienced by individuals (16,46). Tactical practitioners can use these measures periodically to track their personnel. If measures such

Training Load in Tactical Populations

Recovery Scoring Guide

Athletes can fill out this guide over the course of a week to assess their own recovery behaviors. Once they have scored a full day, the total reveals whether they are paying adequate attention to their physical and mental recovery needs. **17-20** daily points is optimal; **15-16** points is good but shows room for improvement; **14** or fewer points means the athlete needs a serious individual evaluation of recovery behaviors.

	Possible Points	Sun.	Mon.	Tues.	Wed.	Thurs.	Fri.	Sat.	SCORING GUIDE ADJUSTMENTS
Nutrition	8								
Breakfast	1								Give ½ a point for a less than full breakfast
Lunch	2								Give one point for a less than full lunch
Dinner	2								Give one point for a less than full dinner
Pre-workout snack	1								
Post-exercise carb refueling within 60 minutes. (recommended: 1.0 to 1.5 g per kilogram of body weight)	2								Give one point for refueling below the recommended amount or for delaying more than 60 minutes.
Hydration	2								
Pre-exercise urine: clear or light color	1								
Post-exercise urine: clear or light color	1								
Sleep and Rest	4								
8 hours of restful sleep	3								Give two points for 7 to <8 hours Give one point for 6-7 hours
Nap during the day	1								
Relaxation and Emotional Status	3								
Fully relaxed 60 minutes post-workout or 30 minutes of feet-up relaxation post workout	1								
No daily psychosocial stress	2								Give one point for mild stress
Stretching/Cooldown	3								
Adequate cooldown after exercise	2								Give one point for partial cooldown
Stretching for at least 10 minutes	1								
TOTALS	20								

Figure 3. TQR Action Scoring Guide. Reproduced from Kentta and Hassmen (46). TQR = Total Quality Recovery.

as these prove to be too time-consuming, implementing muscle soreness and physical fatigue scales may be an alternative method to track IL (64) (Table).

Acute:chronic workload ratio.

Acute:chronic workload ratio (ACWR) is an attempt to provide a more complete picture of an individual's cumulative load by comparing an acute workload with a chronic workload and quantifying the changes over time (26). As mentioned above, it may be the rate of change in load, more than the absolute load, that relates to injury risk (79). The importance of rate of change in load can be historically seen through the 10% rule, which states that increases in exercise frequency, duration, or intensity should be limited to 10% of the previous

week (76). Although the timeframes may vary, in sports the acute workload is typically represented by 1 week, whereas the chronic load is a summation of the previous 3–6 weeks (26). If the acute workload is greater than the chronic workload (signifying increases in TL), the ratio will be greater than 1, and the individual will be in a fatigued state (26). Conversely, if the acute workload is less than chronic workload, the ratio will be less than 1, and injury may be less likely; however, performance may plateau or decline with inadequate stimulus (26). The chronic workload has also been called “fitness” in the literature, whereas the acute workload has been called “fatigue” (26). Evidence shows that high chronic workload or “fitness” may protect against spikes in acute

workload or “fatigue” (26,37,56). In a study by Malone et al. (56), it was found that a high chronic TL (4,750 AU) measured by sRPE protected against injury from maximal velocity running. In another study by Hulin et al. (37), it was found that a high chronic workload (>16,095 m covered) combined with a moderate ACWR (1.03–1.38; relative risk [RR] = 6.2 ± 2.2) resulted in a lower injury risk than low chronic workload (<16,095 meters covered) combined with a moderate ACWR (1.03–1.38; RR = 9.3 ± 2.2). RR states how many times more likely an outcome will occur in an exposed group compared with a nonexposure group (85). In the case of the Hulin et al.'s article (37), those who experienced a high chronic workload were 6.2 times more likely to experience an injury,

Table
Potential examples of load measurements in tactical populations

Method	Example
Internal load	
sRPE	Subjective RPE score × session duration; can be performed after PT sessions, combat training, response calls, and field sessions
HR	Average HR during PT sessions, combat training, response calls, and field sessions
HRV	Each day
TRIMP	After PT sessions, combat training, response calls, and field sessions
Wellness questionnaire	Performed at set time points (daily, weekly, etc.) depending on the questionnaire
External load	
TD	Distance covered during PT sessions, field sessions, and response calls
Speed	Distance covered in various speed bands (10–15, 15–20 km/h, etc.) during physical training and field sessions
Sprints	Number of sprints performed during PT sessions
Volume load	Calculated after strength training sessions (number of repetitions × external load [kg])
Repetition volume	Calculated after strength training sessions (total number of repetitions)

HR = heart rate; HRV = HR variability; PT = Physical Training; RPE = rating of perceived exertion; sRPE = session RPE; TD = total distance.

whereas those in the low chronic workload were 9.3 times more likely to experience an injury. ACWR may be more appropriate to use than the 10% rule, as it is not a rigid law, but can be adjusted based on past training.

One limitation of the ACWR model is that it does not account for the decaying effects of fatigue and fitness over time and therefore may be able to be improved (7,61). A new method using exponential weighted moving averages (EWMAs), instead of rolling averages, was recently put forward by Williams et al. (86). This method attempts to account for the decaying effects by assigning a decreasing weighting for older values (86). A study by Murray et al. (61) compared EWMA ACWR with rolling average ACWR and found that although both models showed a significant increase in injury risk with high ACWR, EWMA was more sensitive to detect increases in injury likelihood at higher ACWRs. When compared with an

ACWR of 1.0–1.49, a rolling average ACWR of 2.0 was shown to have a relative injury risk of 6.52 (4.83–8.80), a statistically significant finding. Using the same comparison, but with EWMA, the RR increased to 21.28 (20.02–22.62)—demonstrating a much higher relative injury risk. Incorporating an EWMA ACWR over a rolling average ACWR may be a more sensitive and advantageous method (86).

Clinicians and practitioners can use ACWR with any of the above-mentioned variables. By incorporating ACWR with measures such as a total distance, sRPE, or volume load, practitioners may be better able to predict how well their personnel will respond to future programs. If knowing that over the past 3 weeks, personnel have only covered an average of 2,000 m per week, practitioners can calculate that covering 4000 m will result in an ACWR of 2.00 and therefore increase injury risk (acute load divided by chronic load, or 4,000 m divided by 2,000 m). Practitioners can adjust their

scheduled training sessions either based on previous work load or based on current occupational demands (e.g., dismounted patrols) to keep personnel within a safe range.

Although ACWR may be a useful tool for predicting injury risk, there are potential limitations affecting its validity. One such issue is the potential spurious correlations that may exist between ACWR and injury risk (52). This is likely due to the fact that the acute load is used in the calculation of the chronic load, as the acute load is on both sides of the equation (52). A second impacting factor is the confounding effect training schedule has on the ability of ACWR to predict injury risk (6). This was calculated with regard to sporting teams, so it remains to be seen if this relationship exists in tactical populations. However, although ACWR may be an effective method to calculate injury risk, it should not be solely relied on by clinicians and practitioners.

The association of training load and injuries.

Research has hypothesized that musculoskeletal injuries may be linked to adverse responses in TL (28), while specifically researched targets regarding TL and injury in the athletic domain have been studied. Research has shown that 2-week cumulative sRPE loads >3,700 AU (32) and high 3-week running distances (73,721–86,662 m) (13) are likely to increase injury risk in Australian Rules football players while studies across multiple sports have shown that ACWR ratios ranging from 0.8 to 1.3 may minimize injury risk while ensuring increases in fitness (22,37). Using either cumulative loads or ACWR provides information to medical staff and strength and conditioning (S&C) coaches, which may assist with reducing injury risk and maximizing performance in tactical environments. It is unlikely that these numbers will be directly transferable given the differences that exist between the tactical and athletic domains, although it should be noted that members of tactical population regularly play sport (1,81).

The associations of training load and fitness and performance.

Although TL may assist with mitigating injury risk, it also is related to fitness and performance. Fitness is a key requirement for tactical personnel and is associated with successful occupational task performance (72), reduced injury risk (48), and mental health benefits (14). However, although training to increase fitness is needed, it may pose a risk if not monitored correctly, as supported by findings that in military personnel, physical training and sports participation are leading causes of injuries (42). Tracking TL may be able to ensure that tactical populations are training at a high enough intensity to increase fitness while not unnecessarily increasing risk of injury (26). In AFL players, it was found that those who had completed high TL (1,600–2,000 AU per week) increased their 2-km time trial

compared with those who completed very high (>2,000 AU per week), moderate (1,250–1,599 AU per week), and low (<1,250 AU per week) TL as measured by sRPE (32). In a study by Gabbett and Domrow (25), it was found that although increases in TL of 175–620 AU, as measured by sRPE, did not result in any increase risk of injury, it did result in decreased performances in agility. These results provide evidence that tracking TL cannot only influence injury risk, but fitness and performance as well. Given the inherent dangers tactical populations can encounter (27,70), decreases in performance or losses of fitness do not simply result in a lost game, but can have life and death consequences. By implementing a method of monitoring TL tactical populations can aim for, and train at a level that maximizes performance gains while minimizing injury risk (7,59).

Although there are notable differences between tactical populations and elite athletes, the TL methodology used in elite sport may provide a template for tactical personnel. Preliminary studies in military populations show the associated increase in injury risk with increase in physical activity (2,83). Although these studies have shown the potential harmful impacts of excessive physical activity, modern methods such as sRPE, ACWR, or any of the other various methods to track EL and IL used in elite sports (7) appear to be suitable to implement to track and adjust TL (26).

Practitioners must recognize that excessive TL is just one potential causative factor of injury in tactical populations. Although there are various other causative factors, that is, smoking (10,48), previous injury (2,10), and poor fitness (48), little research has been performed regarding monitoring TL, especially using the methods seen in elite sport (7,26). The current depth of research is lacking given the impact overloading or underloading has on injuries in tactical populations. Multiple

studies have shown the impact of high physical activity and repetitive microtrauma has on musculoskeletal injuries in tactical populations (33,83). A large amount of these injuries are due to repetitive microtrauma, or overuse, accounting for almost 4 times as many injuries as acute trauma (33). Military recruits especially are at risk of injury due to sudden and large increases in TL due to high amounts of running and other physical activity with possibly little to no previous training (66,77). Given the impact injuries have on both institutional costs, approximately \$1.5 billion (USD) per year in U.S. Army (35,62), and organizational workload, it is vital that tactical populations begin to research and implement these practices.

CONCLUSION

Given both the high physical workload and injury rate within tactical populations, it is important that steps are put into place to maximize performance in decreasing injury risk. A wide range of variables including HR, sRPE, HRV, distance covered, and volume load among others may allow practitioners to monitor TL within their personnel. In addition, ACWRs could be devised to minimize any spikes in TL, which may increase injury risk. A comprehensive understanding of TL may allow for optimizing the programming of activities such as pack marches and patrols, to allow for the addition of or reduction of training where required. For example, if a military unit is planning to complete a 5-km dismounted patrol in 4 weeks, an S&C coach can analyze the distance covered previously during field and physical training sessions. Owing to the minimal research performed within the tactical domain to date, specific ratios are not currently available; however, evidence from elite sport highlights that ACWRs from 0.8 to 1.3 may be optimal for both performance gains and injury reduction.

Conflicts of Interest and Source of Funding: The authors report no conflicts of interest. This article was supported by a Higher Degree Research Scholarship for Mr. Maupin.



Danny Maupin is a PhD student in the Faculty of Health Sciences and Medicine at Bond University.



Ben Schram is an assistant professor in the Faculty of Health Sciences and Medicine at Bond University.



Robin Orr is an associate professor in the Faculty of Health Sciences and Medicine at Bond University.

REFERENCES

- Adams MS and Croft A. Sports injuries in British troops deployed on operation resolute (Bosnia). *J R Army Med Corps* 143: 35, 1997.
- Almeida SA, Williams KM, Shaffer RA, and Brodine SK. Epidemiological patterns of musculoskeletal injuries and physical training. *Med Sci Sports Exerc* 31: 1176–1182, 1999.
- Bannister E. *Modeling Elite Athletic Performance*. Champaign, IL: Human Kinetics, 1991.
- Benson AC, Bruce L, and Gordon BA. Reliability and validity of a GPS-enabled iPhone TM “app” to measure physical activity. *J Sports Sci* 33: 1–8, 2014.
- Borg G. Perceived exertion as an indicator of somatic stress. *Scand J Rehabil Med* 2: 92–98, 1970.
- Bornn L, Ward P, and Norman D. *Training Schedule Confounds the Relationship between Acute:Chronic Workload Ratio and Injury*. Boston, MA: MIT Sloan Sports Analytics Conference, 2019.
- Bourdon PC, Cardinale M, Murray A, Gastin P, Kellmann M, Varley MC, Gabbett TJ, Coutts AJ, Burgess DJ, Gregson W, and Cable NT. Monitoring athlete training loads: Consensus statement. *Int J Sports Physiol Perform* 12(Suppl 2): S2161–S170, 2017.
- Bovens A, Janssen G, Vermeer H, Hoeberigs J, Janssen M, and Verstappen F. Occurrence of running injuries in adults following a supervised-training program. *Int J Sports Med* 10: S186–S190, 1989.
- Brandl SG and Strohshine MS. The physical hazards of police work revisited. *Police Q* 15: 262–282, 2012.
- Bulzacchelli MT, Sulsky SI, Rodriguez-Monguio R, Karlsson LH, and Hill MOT. Injury during U.S. Army basic combat training: A systematic review of risk factor studies. *Am J Prev Med* 47: 813–822, 2014.
- Carbone PD, Carlton SD, Stierli M, and Orr R. The impact of load carriage on the marksmanship of the tactical police officer: A pilot study. *J Aust Strength Cond* 22: 50–57, 2014.
- Carlton A, Gorey R, and Orr R. The impact of suppressing a structural fire on firefighter hydration. *J Aust Strength Cond* 24: 27–33, 2016.
- Colby MJ, Dawson B, Heasman J, Rogalski B, and Gabbett TJ. Accelerometer and GPS-derived running loads and injury risk in elite Australian footballers. *J Strength Cond Res* 28: 2244–2252, 2014.
- Crowley SK, Wilkinson LL, Wigfall LT, Reynolds AM, Muraca ST, Glover SH, Wooten NR, Sui X, Beets MW, Durstine JL, Newman-Norlund RD, and Youngstedt SD. Physical fitness and depressive symptoms during army basic combat training. *Med Sci Sports Exerc* 47: 151–158, 2015.
- Cummins C, Orr R, O'Connor H, and West C. Global positioning systems (GPS) and microtechnology sensors in team sports: A systematic review. *Sports Med* 43: 1025–1042, 2013.
- Davis H, Orzeck T, and Keelan P. Psychometric item evaluations of the Recovery-Stress Questionnaire for athletes. *Psychol Sport Exerc* 8: 917–938, 2007.
- Dawes JJ, Orr RM, Flores RR, Lockie RG, Kornhauser C, and Holmes R. A physical fitness profile of state highway patrol officers by gender and age. *Ann Occup Environ Med* 29, 2017.
- DiVencenzo HR, Morgan AL, Laurent CM, and Keylock KT. Metabolic demands of law enforcement personal protective equipment during exercise tasks. *Ergonomics* 57: 1760–1765, 2014.
- Drain J, Orr R, Attwells R, and Billing D. *Load Carriage Capacity of the Dismounted Combatant-A Commanders' Guide*. Victoria, Australia: Defence Science and Technology Organisation, Human Protection and Performance Division, 2012.
- Eckard TG, Padua DA, Hearn DW, Pexa BS, and Frank BS. The relationship between training load and injury in athletes: A systematic review. *Sports Med* 48: 1–33, 2018.
- Edwards S. High performance training and racing. In: *The Heart Rate Monitor Book*. Sacramento, CA: Fleet Feet Press, 1994. pp. 113–123.
- Ehrmann FE, Duncan CS, Sindhusake D, Franzsen WN, and Greene DA. GPS and injury prevention in professional soccer. *J Strength Cond Res* 30: 360–367, 2016.
- Farrell PA, Wilmore J, Coyle E, Billing J, and Costill DL. Plasma lactate accumulation and distance running performance. *Med Sci Sports Exerc* 25: 1091–1097, 1993.
- Foster C, Florhaug JA, Franklin J, Gottschall L, Hrovatin LA, Parker S, Doleshal P, and Dodge C. A new approach to monitoring exercise training. *J Strength Cond Res* 15: 109–115, 2001.
- Gabbett TJ and Domrow N. Relationships between training load, injury, and fitness in sub-elite collision sport athletes. *J Sports Sci* 25: 1507–1519, 2007.
- Gabbett TJ. The training-injury prevention paradox: Should athletes be training smarter and harder? *Br J Sports Med* 50: 273–280, 2016.
- Galdorisi G. The 2016 Australian defence white paper: A view from across the pacific. *Asia Pac Def Rep* 42: 34–38, 2016.
- Gisselman AS, Baxter GD, Wright A, Hegedus E, and Tumilty S. Musculoskeletal overuse injuries and heart rate variability: Is there a link? *Med Hypotheses* 87: 1–7, 2016.
- Haff GG and Triplett NT. *Essentials of Strength Training and Conditioning* (4th ed). Champaign, IL: Human Kinetics, 2015.
- Halson SL. Monitoring training load to understand fatigue in athletes. *Sports Med* 44(Suppl 2): S139–S147, 2014.
- Hamill RW, Shapiro RE, and Vizzard MA. Chapter 4—Peripheral autonomic nervous system. In: *Primer on the Autonomic Nervous System* (3rd Ed). Robertson D, Biaggioni I, Burnstock G, Low PA, and

Training Load in Tactical Populations

- Paton JFR, eds. San Diego, CA: Academic Press, 2012. pp. 17–26.
32. Harrison PW and Johnston RD. Relationship between training load, fitness, and injury over an Australian rules football preseason. *J Strength Cond Res* 31: 2686–2693, 2017.
 33. Hauschild V, Hauret K, Richardson M, Jones BH, and Lee T. *A Taxonomy of Injuries for Public Health Monitoring and Reporting*. Gunpowder, MD: Army Public Health Center, Aberdeen Proving Ground, 2017.
 34. Hedelin R, Wiklund U, Bjerle P, and Henriksson-Larsén K. Cardiac autonomic imbalance in an overtrained athlete. *Med Sci Sports Exerc* 32: 1531–1533, 2000.
 35. Hilyer JC, Brown KC, Sirls AT, and Peoples L. A flexibility intervention to reduce the incidence and severity of joint injuries among municipal firefighters. *J Occup Med* 32: 631–637, 1990.
 36. Hoyt RW, Opstad PK, Haugen A-H, DeLany JP, Cymerman A, and Friedl KE. Negative energy balance in male and female rangers: Effects of 7 d of sustained exercise and food deprivation. *Am J Clin Nutr* 83: 1068–1075, 2006.
 37. Hulin BT, Gabbett TJ, Lawson DW, Caputi P, and Sampson JA. The acute: Chronic workload ratio predicts injury: High chronic workload may decrease injury risk in elite rugby league players. *Br J Sports Med* 50: 231–236, 2015.
 38. Hunt AP, Orr RM, and Billing DC. Developing physical capability standards that are predictive of success on Special Forces selection courses. *Mil Med* 178: 619–624, 2013.
 39. Impellizzeri FM, Rampinini E, Coutts AJ, Sassi A, and Marcora SM. Use of RPE-based training load in soccer. *Med Sci Sports Exerc* 36: 1042–1047, 2004.
 40. Johnston RJ, Watsford ML, Kelly SJ, Pine MJ, and Spurr RW. Validity and interunit reliability of 10 Hz and 15 Hz GPS units for assessing athlete movement demands. *J Strength Cond Res* 28: 1649–1655, 2014.
 41. Johnston R, Cahalan R, O’Keeffe M, O’Sullivan K, and Comyns T. The associations between training load and baseline characteristics on musculoskeletal injury and pain in endurance sport populations: A systematic review. *J Sci Med Sport* 21: 910–918, 2018.
 42. Jones B and Knapik J. Physical training and exercise-related injuries. *Sports Med* 27: 111–125, 1999.
 43. Kaufman KR, Brodine S, and Shaffer R. Military training-related injuries: Surveillance, research, and prevention. *Am J Prev Med* 18: 54–63, 2000.
 44. Keeler JM. *The Effect of Tactical Tasks and Gear on Muscle Activation of SWAT Officers [master of science]*. Kinesiology and Health Promotion, University of Kentucky, Lexington KY, 2014, Theses and Dissertations.
 45. Kellmann M and Günther KD. Changes in stress and recovery in elite rowers during preparation for the Olympic Games. *Med Sci Sports Exerc* 32: 676–683, 2000.
 46. Kenttä G and Hassmén P. Overtraining and recovery: A conceptual model. *Sports Med* 26: 1–16, 1998.
 47. Kenttä G and Hassmén P. *Train Smart: Avoid Overtraining Syndrome*. Stockholm, Sweden: SISU Idrottsböcker, 1999.
 48. Knapik JJ, Sharp MA, Canham-Chervak M, Hauret K, Patton JF, and Jones BH. Risk factors for training-related injuries among men and women in basic combat training. *Med Sci Sports Exerc* 33: 946–954, 2001.
 49. Knapik JJ, Reynolds KL, and Harman E. Soldier load carriage: Historical, physiological, biomechanical, and medical aspects. *Mil Med* 169: 45–56, 2004.
 50. Le Meur Y, Pichon A, Schaal K, Schmitt L, Louis J, Gueneron J, Vidal PP, and Hausswirth C. Evidence of parasympathetic hyperactivity in functionally overreached athletes. *Med Sci Sports Exerc* 45: 2061–2071, 2013.
 51. Lisman P, Connor FG, Deuster PA, and Knapik JJ. Functional movement screen and aerobic fitness predict injuries in military training. *Med Sci Sports Exerc* 45: 636, 2013.
 52. Lolli L, Batterham AM, Hawkins R, Kelly DM, Strudwick AJ, Thorpe R, Gregson W, and Atkinson G. Mathematical coupling causes spurious correlation within the conventional acute-to-chronic workload ratio calculations. *Br J Sports Med*, 2017 [Epub ahead of print].
 53. Lyons K, Radburn C, Orr R, and Pope R. A profile of injuries sustained by law enforcement officers: A critical review. *Int J Environ Res Public Health* 14: 142–162, 2017.
 54. Lysholm J and Wiklander J. Injuries in runners. *Am J Sports Med* 15: 168–171, 1987.
 55. Malone S and Collins K. Relationship between individualized training impulse and aerobic fitness measures in hurling players across a training period. *J Strength Cond Res* 30: 3140–3145, 2016.
 56. Malone S, Roe M, Doran DA, Gabbett TJ, and Collins K. High chronic training loads and exposure to bouts of maximal velocity running reduce injury risk in elite Gaelic football. *J Sci Med Sport* 20: 250–254, 2017.
 57. Manzi V, Iellamo F, Impellizzeri F, D’Ottavio S, and Castagna C. Relation between individualized training impulses and performance in distance runners. *Med Sci Sports Exerc* 41: 2090–2096, 2009.
 58. Mohr M, Krstrup P, and Bangsbo J. Match performance of high-standard soccer players with special reference to development of fatigue. *J Sports Sci* 21: 519–528, 2003.
 59. Morton RH. Modeling training and overtraining. *J Sports Sci* 15: 335–340, 1997.
 60. Mouro L, Bouhaddi M, Perrey S, Cappelle S, Henriot MT, Wolf JP, Rouillon JD, and Regnard J. Decrease in heart rate variability with overtraining: Assessment by the Poincaré plot analysis. *Clin Physiol Funct Imaging* 24: 10–18, 2004.
 61. Murray NB, Gabbett TJ, Townshend AD, and Blanch P. Calculating acute: Chronic workload ratios using exponentially weighted moving averages provides a more sensitive indicator of injury likelihood than rolling averages. *Br J Sports Med* 51: 749–754, 2017.
 62. Nindl BC, Williams TJ, Deuster PA, Butler NL, and Jones BH. Strategies for optimizing military physical readiness and preventing musculoskeletal injuries in the 21st century. *US Army Med Dep J* 8: 5–23, 2013.
 63. Ogle AD and Young JA. USAF special tactics operator combat exposure and psychological stress. *Mil Psychol* 28: 123–133, 2016.
 64. O’Leary TJ, Saunders SC, McGuire SJ, Venables MC, and Izard RM. Sex differences in training loads during British army basic training. *Med Sci Sports Exerc* 50: 2565–2574, 2018.
 65. Orr R, Pope R, Coyle J, and Johnston V. Occupational loads carried by Australian soldiers on military operations. *J Health Saf Environ* 31: 451–467, 2015.
 66. Orr R, Ford K, and Stierli M. Implementation of an ability-based training program in police force recruits. *J Strength Cond Res* 30: 2781–2787, 2016.
 67. Orr R, Pope R, Johnston V, and Coyle J. Soldier load carriage: An investigation into the load carriage conditioning practices of the Australian Regular Army. Paper presented at: Australian Military Medicine Association; October 13, 2012; Brisbane, Australia.

68. Piggott B. *The Relationship between Training Load and Incidence of Injury and Illness over a Pre-season at an Australian Football League Club*: Faculty of Computing, Health and Science, Edith Cowan University, Joondalup WA, Australia, 2008.
69. Pope R, Orr R, and MacDonald D. Incidence rates of reported work health and safety incidents and injuries in part-time and full-time Australian Army personnel. *J Mil Veterans Health* 23: 45–46, 2015.
70. Pryor RR, Colburn D, Crill MT, Hostler DP, and Suyama J. Fitness characteristics of a suburban special weapons and tactics team. *J Strength Cond Res* 26: 752–757, 2012.
71. Robinson M, Siddall A, Bilzon J, Thompson D, Greeves J, Izard R, and Stokes K. Low fitness, low body mass and prior injury predict injury risk during military recruit training: A prospective cohort study in the British army. *BMJ Open Sport Exerc Med* 2: e000100, 2016.
72. Robinson J, Roberts A, Irving S, and Orr R. Aerobic fitness is of greater importance than strength and power in the load carriage performance of specialist police. *Int J Exerc Sci* 11: 987–998, 2018.
73. Rossi R. Fire fighting and its influence on the body. *Ergonomics* 46: 1017–1033, 2003.
74. Sallis JF and Saelens BE. Assessment of physical activity by self-report: Status, limitations, and future directions. *Res Q Exerc Sport* 71: 1–14, 2000.
75. Scott TJ, Black CR, Quinn J, and Coutts AJ. Validity and reliability of the session-RPE method for quantifying training in Australian football: A comparison of the CR10 and CR100 scales. *J Strength Cond Res* 27: 270–276, 2013.
76. Sheppard MJ and Triplett NT. *Essentials of Strength Training and Conditioning* (4th ed). Champaign, IL: Human Kinetics, 2016.
77. Sherrard J, Lenné M, Cassell E, Stokes M, and Ozanne-Smith J. Injury prevention during physical activity in the Australian Defence Force. *J Sci Med Sport* 7: 106–117, 2004.
78. Simpson RJ, Graham SM, Connaboy C, Clement R, Pollonini L, and Florida-James GD. Blood lactate thresholds and walking/running economy are determinants of backpack-running performance in trained soldiers. *Appl Ergon* 58: 566–572, 2017.
79. Soligard T, Schweltnus M, Alonso JM, Bahr R, Clarsen B, Dijkstra HP, Gabbett T, Gleeson M, Häggglund M, Hutchinson MR, Janse van Rensburg C, Khan KM, Meeusen R, Orchard JW, Pluim BM, Raftery M, Budgett R, and Engebretsen L. How much is too much? (Part 1) International Olympic Committee consensus statement on load in sport and risk of injury. *Br J Sports Med* 50: 1030–1041, 2016.
80. Strath SJ, Swartz AM, Bassett DR, O'Brien WL, King GA, and Ainsworth BE. Evaluation of heart rate as a method for assessing moderate intensity physical activity. *Med Sci Sports Exerc* 32(9 Suppl): S465, 2000.
81. Strowbridge N and Burgess K. Sports and training injuries in British Soldiers: The Colchester Garrison Sports Injury and Rehabilitation Centre. *J R Army Med Corps* 148: 236, 2002.
82. Taylor NA, Dodd MJ, Taylor EA, and Donohoe AM. A retrospective evaluation of injuries to Australian urban firefighters (2003 to 2012): Injury types, locations, and causal mechanisms. *J Occup Environ Med* 57: 757–764, 2015.
83. Trank TV, Ryman DH, Minagawa RY, Trone DW, and Shaffer RA. Running mileage, movement mileage, and fitness in male U.S. Navy recruits. *Med Sci Sports Exerc* 33: 1033–1038, 2001.
84. Veenstra BJ, Wyss T, Roos L, Delves SK, Buller M, and Beeler N. An evaluation of measurement systems estimating gait speed during a loaded military march over graded terrain. *Gait Posture* 61: 204–209, 2018.
85. Viera AJ. Odds ratios and risk ratios: what's the difference and why does it matter? *South Med J* 101: 730, 2008.
86. Williams S, West S, Cross MJ, and Stokes KA. Better way to determine the acute: chronic workload ratio? *Br J Sports Med* 51: 209–210, 2017.